

A short proof of Cayley's Tree Formula

Alok B. Shukla
Nashik, India
sajal.eee@gmail.com

October 9, 2009

Abstract

A simple combinatorial argument employing symmetry of edges is put forth to establish a recursive relation for counting the number of different labeled trees T_n on n vertices.

$$T_n = \frac{n}{2} \sum_{k=0}^{n-2} \binom{n-2}{k} T_{k+1} T_{n-k-1}; \text{ for } n > 1 \text{ and } T_1 = 1$$

It is proved that T_n calculated from this recursive relation is equivalent to the well known Cayley's Tree Formula n^{n-2} .

Theorem: *There are exactly $(n)^{n-2}$ labeled trees on n vertices for each $n \geq 1$.*

Proof: Let $\mathbf{T}$ represent the set of all labeled trees on $\mathbf{n}$ vertices of given vertex set $\mathbf{V}\{\mathbf{v}_1, \mathbf{v}_2 \dots \mathbf{v}_n\}$. Let $\mathbf{T}_n$ denote the cardinality of $\mathbf{T}$. Also let $\mathbf{E}_n$ denote the number of trees in $\mathbf{T}$ that always contain one specified edge, say $\mathbf{v}_1 \mathbf{v}_2$.

It should be noted that $\mathbf{T}$ is not affected by any rearrangement of vertices of $\mathbf{V}$. This symmetry implies that all the edges are equivalent and contribute equally to $\mathbf{T}$. As the number of possible edges is $\binom{n}{2} = \frac{1}{2}n(n-1)$; the total number of edges in all trees in $\mathbf{T}$ equals

$$\frac{1}{2} n (n-1) \mathbf{E}_n = (n-1) \mathbf{T}_n$$

where the right side of equality follows from the fact that each one of $\mathbf{T}_n$ trees in $\mathbf{T}$ contains $(n-1)$ edges. Therefore we have for all $n > 1$

$$\mathbf{T}_n = \frac{1}{2} n \mathbf{E}_n \tag{1}$$

Further, it is easy to see that $\mathbf{T}_1 = 1$.

Without any loss of generality, let us fix edge $\mathbf{v}_1 \mathbf{v}_2$ and count the number of possible trees to determine $\mathbf{E}_n$. For this, k vertices are chosen from the set

$V\{v_3, v_4, \dots, v_n\}$ in $\binom{n-2}{k}$ ways. These k vertices together with vertex v_1 form a set of $(k+1)$ vertices and would give T_{k+1} trees. The remaining $(n-2-k)$ vertices are attached to vertex v_2 to form a set of $(n-k-1)$ vertices to yield T_{n-k-1} trees. By summing for $k=0$ to $n-2$ we get,

$$E_n = \sum_{k=0}^{n-2} \binom{n-2}{k} T_{k+1} T_{n-k-1} \quad (2)$$

From Eq(1) and Eq(2) we get

$$T_n = \frac{1}{2} n \sum_{k=0}^{n-2} \binom{n-2}{k} T_{k+1} T_{n-k-1} \quad (3)$$




Figure 1: Counting all Trees containing edge V_1V_2

Let the exponential generating function $T(S)$ be defined as

$$T(S) = \sum_{n=1}^{\infty} T_n \frac{S^n}{(n-1)!} \quad (4)$$

$$\begin{aligned} (T(S))^2 &= \sum_{n=1}^{\infty} \sum_{k=0}^{n-2} \left\{ T_{k+1} \frac{S^{k+1}}{(k!)} \right\} \left\{ T_{n-k-1} \frac{S^{n-k-1}}{(n-k-2)!} \right\} \\ &= \sum_{n=1}^{\infty} \sum_{k=0}^{n-2} \left\{ \binom{n-2}{k} T_{k+1} T_{n-k-1} \frac{S^n}{(n-2)!} \right\} \end{aligned}$$

By using the value of T_n from the recurrence relation of Eq (3),

$$(T(S))^2 = \sum_{n=1}^{\infty} \left\{ 2 \frac{(n-1)}{n} T_n \frac{S^n}{(n-1)!} \right\} \quad (5)$$

On differentiating with respect to $\mathbf{S}$,

$$2\mathbf{T}(\mathbf{S})\mathbf{T}'(\mathbf{S}) = \sum_{\mathbf{n}=1}^{\infty} \left\{ 2(\mathbf{n}-1)\mathbf{T}_{\mathbf{n}} \frac{\mathbf{S}^{\mathbf{n}-1}}{(\mathbf{n}-1)!} \right\}$$

$$\mathbf{T}(\mathbf{S})\mathbf{T}'(\mathbf{S}) = \sum_{\mathbf{n}=1}^{\infty} \left\{ \mathbf{n}\mathbf{T}_{\mathbf{n}} \frac{\mathbf{S}^{\mathbf{n}-1}}{(\mathbf{n}-1)!} \right\} - \frac{1}{\mathbf{S}} \sum_{\mathbf{n}=1}^{\infty} \left\{ \mathbf{T}_{\mathbf{n}} \frac{\mathbf{S}^{\mathbf{n}}}{(\mathbf{n}-1)!} \right\} = \mathbf{T}'(\mathbf{S}) - \frac{\mathbf{T}(\mathbf{S})}{\mathbf{S}}$$

$$\partial[\mathbf{T}(\mathbf{S})] = \partial[\ln \left[\frac{\mathbf{T}(\mathbf{S})}{\mathbf{S}} \right]]$$

On integrating,

$$\mathbf{T}(\mathbf{S}) = \ln \left[\frac{\mathbf{T}(\mathbf{S})}{\mathbf{S}} \right] + \mathbf{C} \quad (6)$$

Where $\mathbf{C}$ is a constant of integration.

To fix $\mathbf{C}$, put $\mathbf{S}=0$ in Eq(6) and note that from Eq(4) - $T(0) = 0$ and $\frac{T(\mathbf{S})}{\mathbf{S}}|_{\mathbf{S}=0} = T_1 = 1$. So we have $0 = \ln(1) + \mathbf{C} \Rightarrow \mathbf{C} = 0$ And Eq(6) reduces to

$$\mathbf{T}(\mathbf{S}) = \mathbf{S} \exp^{\mathbf{T}(\mathbf{S})} \quad (7)$$

One can use **Lagrange Inversion Theorem** to expand $T(S)$ in power series of S and the result is

$$\mathbf{T}(\mathbf{S}) = \sum_{\mathbf{n}=1}^{\infty} \frac{\mathbf{n}^{\mathbf{n}-1}}{\mathbf{n}!} \mathbf{S}^{\mathbf{n}} = \sum_{\mathbf{n}=1}^{\infty} \mathbf{n}^{\mathbf{n}-2} \frac{\mathbf{S}^{\mathbf{n}}}{(\mathbf{n}-1)!} \quad (8)$$

By equating the coefficient of S^n in Eq (4) and Eq (8) we get $\mathbf{T}_{\mathbf{n}} = \mathbf{n}^{\mathbf{n}-2}$. This completes the proof.

References

- [1] John M. Harris , Jeffry L. Hirst and Michael J. Mossinghoff *Combinatorics and Graph Theory*, Second Edition, 2008 Springer Publication
- [2] Philippe Flajolet and Robert Sedgewick *Analytic Combinatorics*, 2009 Cambridge University Press
- [3] Herbert S. Wilf *generatingfunctionology*, 1994 Academic Press, Inc